

ISSN 2518-1467 (Online),
ISSN 1991-3494 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Ш Ы С Ы

ВЕСТНИК

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

THE BULLETIN

THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

PUBLISHED SINCE 1944

4

JULY – AUGUST 2020

ALMATY, NAS RK

NAS RK is pleased to announce that Bulletin of NAS RK scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of Bulletin of NAS RK in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential multidiscipline content to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабаршысы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабаршысының Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді мультидисциплинарлы контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Вестник НАН РК» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Вестника НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному мультидисциплинарному контенту для нашего сообщества.

Б а с р е д а к т о р ы

х.ғ.д., проф., ҚР ҰҒА академигі

М.Ж. Жұрынов

Р е д а к ц и я а л қ а с ы:

Абиев Р.Ш. проф. (Ресей)
Абылкасымова А.Е. проф., академик (Қазақстан)
Аврамов К.В. проф. (Украина)
Аппель Юрген проф. (Германия)
Баймұқанов Д.А. проф., академик (Қазақстан)
Баймұратов У.Б. проф., академик (Қазақстан)
Байтанаев Б.А. проф., академик (Қазақстан)
Байтулин И.О. проф., академик (Қазақстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Қазақстан)
Велесько С. проф. (Германия)
Велихов Е.П. проф., РҒА академигі (Ресей)
Кабульдинов З.Е. проф. (Қазақстан)
Қажыбек Е.З. проф., корр.-мүшесі (Қазақстан)
Қалимолдаев М.Н. проф., академик (Қазақстан), бас ред. орынбасары
Қамзабекұлы Д. проф., академик (Қазақстан)
Қойгелдиев М.К. проф., академик (Қазақстан)
Лупашку Ф. проф., корр.-мүшесі (Молдова)
Мохд Хасан Селамат проф. (Малайзия)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., корр.-мүшесі (Қазақстан)
Полещук О.Х. проф. (Ресей)
Поняев А.И. проф. (Ресей)
Сагиян А.С. проф., академик (Армения)
Таймагамбетов Ж.К. проф., академик (Қазақстан)
Хрипунов Г.С. проф. (Украина)
Шәукенова З.К. проф., корр.-мүшесі (Қазақстан)
Юлдашбаев Ю.А. проф., РҒА академигі (Ресей)
Якубова М.М. проф., академик (Тәжікстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабаршысы».

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.).

Қазақстан Республикасының Ақпарат және коммуникациялар министрлігінің Ақпарат комитетінде 12.02.2018 ж. берілген № 16895-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://www.bulletin-science.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Г л а в н ы й р е д а к т о р
д.х.н., проф. академик НАН РК
М.Ж. Журинов

Р е д а к ц и о н н а я к о л л е г и я:

Абиев Р.Ш. проф. (Россия)
Абылкасымова А.Е. проф., академик (Казахстан)
Аврамов К.В. проф. (Украина)
Апель Юрген проф. (Германия)
Баймуканов Д.А. проф., академик (Казахстан)
Баймуратов У.Б. проф., академик (Казахстан)
Байтанаев Б.А. проф., академик (Казахстан)
Байтулин И.О. проф., академик (Казахстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Казахстан)
Велесько С. проф. (Германия)
Велихов Е.П. проф., академик РАН (Россия)
Кабульдинов З.Е. проф. (Казахстан)
Кажыбек Е.З. проф., чл.-корр. (Казахстан)
Калимолдаев М.Н. академик (Казахстан), зам. гл. ред.
Камзабекулы Д. проф., академик (Казахстан)
Койгельдиев М.К. проф., академик (Казахстан)
Лунашку Ф. проф., чл.-корр. (Молдова)
Мохд Хасан Селамат проф. (Малайзия)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., чл.-корр. (Казахстан)
Полещук О.Х. проф. (Россия)
Поняев А.И. проф. (Россия)
Сагиян А.С. проф., академик (Армения)
Таймагамбетов Ж.К. проф., академик (Казахстан)
Хрипунов Г.С. проф. (Украина)
Шаукенова З.К. проф., чл.-корр. (Казахстан)
Юлдашбаев Ю.А. проф., академик РАН (Россия)
Якубова М.М. проф., академик (Таджикистан)

«Вестник Национальной академии наук Республики Казахстан».

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и коммуникаций и Республики Казахстан № 16895-Ж, выданное 12.02.2018 г.

Периодичность: 6 раз в год.

Тираж: 2000 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18.

<http://www.bulletin-science.kz/index.php/en/>

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNazGRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief

doctor of chemistry, professor, academician of NAS RK

M.Zh. Zhurinov

Editorial board:

Abiyev R.Sh. prof. (Russia)
Abylkasymova A.E. prof., academician (Kazakhstan)
Avramov K.V. prof. (Ukraine)
Appel Jurgen, prof. (Germany)
Baimukanov D.A. prof., academician (Kazakhstan)
Baimuratov U.B. prof., academician (Kazakhstan)
Baitanaev B.A. prof., academician (Kazakhstan)
Baitullin I.O. prof., academician (Kazakhstan)
Joseph Banas, prof. (Poland)
Bersimbayev R.I. prof., academician (Kazakhstan)
Velesco S., prof. (Germany)
Velikhov Ye.P. prof., academician of RAS (Russia)
Kabuldinov Z.E. prof. (Kazakhstan)
Kazhybek E.Z. prof., corr. member. (Kazakhstan)
Kalimoldayev M.N. prof., academician (Kazakhstan), deputy editor in chief
Kamzabekuly D. prof., academician (Kazakhstan)
Koigeldiev M.K. prof., academician (Kazakhstan)
Lupashku F. prof., corr. member (Moldova)
Mohd Hassan Selamat, prof. (Malaysia)
Nowak Isabella, prof. (Poland)
Ogar N.P. prof., corr. member (Kazakhstan)
Poleshchuk O.Kh. prof. (Russia)
Ponyaev A.I. prof. (Russia)
Sagiyani A.S. prof., academician (Armenia)
Tajmagambetov Zh.K. prof., academician (Kazakhstan)
Khripunov G.S. prof. (Ukraine)
Shaukenova Z.K. prof., corr. member. (Kazakhstan)
Yuldashbayev Y.A., prof., academician of RAS (Russia)
Yakubova M.M. prof., academician (Tadjikistan)

Bulletin of the National Academy of Sciences of the Republic of Kazakhstan.

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Communications of the Republic of Kazakhstan No. **16895-Ж**, issued on 12.02.2018.

Periodicity: 6 times a year.

Circulation: 2000 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://www.bulletin-science.kz/index.php/en/>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

**N. N. Kuzmina¹, O. Yu. Petrov¹, V. G. Semenov²,
D. A. Baimukanov³, A. I. Lyubimov⁴, Kh. A. Aubakirov⁵**

¹Mari State University, Yoshkar-Ola, Mari El Republic, Russia;

²Chuvash State Agricultural Academy, Cheboksary, Chuvash Republic, Russia;

³Non-Commercial Joint-Stock Company «Kazakh National Agrarian University», Almaty, Kazakhstan;

⁴Izhevsk State Agricultural Academy, Izhevsk, Udmurt Republic, Russia;

⁵M. H. Dulati Taraz State University, Taraz, Kazakhstan.

E-mail: kuzmina221995@mail.ru, tmspetrov@yandex.ru, semenov_v.g@list.ru,
dbaimukanov@mail.ru, rector@izhgsha.ru, hamit_a57@mail.ru

STUDY OF THE EFFICIENCY OF NATURAL ANTIOXIDANTS IN STORING POULTRY RAW MATERIAL PROCESSING

Abstract. The efficiency of using dihydroquercetin as an antioxidant, which increases the duration of storage of poultry processing raw materials, is investigated.

A comparative assessment of the effectiveness of the natural antioxidants of vitamins E, C, rutin, and dihydroquercetin was carried out for the duration of storage of minced meat from the fillet, meat of deboned meat and skin from broiler carcasses. The introduction of antioxidants in the experimental samples of raw materials ensured, during 28 days of storage, a lower value of the indicators of oxidative spoilage: the acid number by an average of 74%, and the peroxide number by 1.5-1.7 times, relative to these indicators in the control sample antioxidants.

The use of dihydroquercetin in experimental samples of frozen mechanically deboned meat for long-term storage, in the studied concentrations, ensured after 1 month of storage a lower value of the oxidative spoilage indicators: the acid number was 2–10% lower, and the peroxide number was 1.5 and 1.7 times lower, relative to these indicators in the control sample.

The results confirm the high efficiency of the use of dihydroquercetin as an active antioxidant, which makes it possible to use it along with existing analogues.

Key words: poultry processing, skin from broiler chicken carcasses, mechanically deboned meat, filet of chilled broiler chicken carcasses, natural antioxidants, vitamins E, C, rutin, dihydroquercetin, oxidation products, storage life.

Introduction. Currently, in the meat industry, the creation and production with a functional orientation with a prolonged shelf life are actual. The consumption of such products allows us to stabilize the metabolism in the body and to improve human health. This is an important aspect for the consumer, and the manufacturer, at the same time, gets the opportunity to produce products of guaranteed quality, taking into account unforeseen situations associated with the deviation of temperature conditions of storage, transportation, and sale [1]. But one of the problems in the production of food, in particular, meat, is the extension of the shelf life while maintaining their quality.

The processes of fat oxidation have a negative effect not only on food, but also on the human body, the most dangerous being the appearance and accumulation of free radicals that contribute to the emergence of a number of specific diseases [2]. The ability of dihydroquercetin to bind and intercept such radicals prevents the development of these diseases.

Most modern synthetic antioxidants and their effects on the human body are not well studied, which leads to ambiguous consequences, and their use is inappropriate since the body has its own antioxidant system. In the body, this function is performed by enzymes (catalase, superoxide dismutase, peroxidase, glutathione peroxidase, ubiquinone), albumin and a number of other proteins, including sulfur-containing and selenium-containing, high-density lipoproteins, steroid hormones, bilirubin, uric acid. These substances are helped by exogenous antioxidants - vitamins that enter the body with food. Therefore, one

of the modern trends in meat production for healthy nutrition is the orientation to the use of natural food additives, which can have a pronounced positive effect on the human body. In this regard, safe natural antioxidants deserve much attention, which not only significantly inhibit the oxidation in meat products during storage, but also serve as the active principle of therapeutic and prophylactic products, which makes their use relevant in formulations of a wide range of products.

Dihydroquercetin is one of the most active antioxidants of natural origin used in food production; recognized dihydroquercetin is a natural compound isolated from Siberian larch wood [3]. Russia, in turn, is the only country in the world that has its unique natural resources in volumes suitable for industrial and cost-effective production.

Dihydroquercetin has the highest activity. Antioxidant activity is an indicator that reflects the ability to inactivate free oxygen radicals. The antioxidant activity of dihydroquercetin manifests itself at its concentrations of 10^{-4} - 10^{-5} . This is the lowest concentration of a substance with antioxidant properties relative to all known exogenous antioxidants, including vitamins A, B, C, D, E, K, β -carotene [4,5].

Adding of dihydroquercetin provides food with additional competitive advantages [4,6].

In addition, the use of natural antioxidants in the production of meat products is due to certain difficulties at the stages of the technological process. Natural antioxidant vitamins easily lose their activity and their structure is destroyed under the influence of various factors: contact with a metal surface, exposure to open sunlight and oxygen, high-temperature treatment (except vitamins A, E, K, and carotenoids) [4,5].

Research conducted at the Sechenov Moscow Medical Academy confirmed that dihydroquercetin is non-toxic, physiologically harmless to human health, does not impart any taste or smell to extraneous products, does not change their color when used. The additive is resistant to temperature (from minus 50 to plus 180 °C), mechanical stresses, and processes occurring in the manufacture of products, that is, it meets all the requirements for whole food additives and, in particular, antioxidants. This compound is included in the list of food additives that do not have a harmful effect on human health when used for cooking food (SanPiN 2.3.2.1078-01 "Hygienic requirements for food safety and nutritional value") [7].

Conducted by relevant Russian institutes (All-Russian Scientific Research Institute of the Dairy Industry, Scientific Research Institute of Storage Problems of the Federal Agency for State Reserves, Kemerovo Technological Institute of Food Industry, All-Russian Research Institute of Meat Industry named after V.M. Gorbатов, All-Russian Scientific Research Institute of Butter making and Cheesemaking, etc.), the studies allowed to develop new technologies and products using dihydroquercetin as a natural antioxidant. Its recommended volume, depending on the fatty acid composition, is 0.005-0.03% by weight of the canned product.

Methods of research. In the first series of studies, during 28 days of storage in a cooled state, in laboratory conditions, comparative studies of the most common natural antioxidants were conducted according to their ability to inhibit the formation of lipid fraction oxidation products in experimental samples of raw materials used in the production of poultry meat products. Antioxidants were added to the experimental finely divided, homogenized samples of raw materials, in accordance with the recommended dosage indicated in table 1.

Table 1 – Dosage options for natural antioxidants in experimental samples of poultry processing raw materials

Antioxidant	Rate of hydration	Raw material		
		fillet	Mechanically deboned meat	Skin
Rutin, mg/kg	1:3	0.59	0.57	0.56
Vitamin C, mg/kg	1:2	0.57	0.56	0.53
Vitamin E, mg/kg	–	0.57	0.56	0.52
Dihydroquercetin, mg/kg	1:3	0.62	0.58	0.57

After 7, 14, 21, and 28 days of storage at a temperature of 3 ± 1 °C, there was a determination in a 3-fold repetition of the content of oxidation products in the control and experimental samples according to acid and peroxide numbers, by common methods.

The use of these antioxidants is regulated by the Methodological Recommendations of the State Sanitary and Epidemiological Regulation of the Russian Federation No. 2.3.1.1915-04 of 2004 "Recommended levels of consumption of food and biologically active substances", establishing adequate and upper acceptable levels of dihydroquercetin (DHQ) consumption in the amount of 25 and 100 mg per

day, vitamin C - 70 and 700 mg per day, vitamin E - 15 and 100 mg per day, rutin - 30 (transferred to rutin) and 100 mg (transferred to rutin), in the food industry [7,8].

Antioxidant preparations, with the exception of vitamin E, were hydrated before being added to the samples for more uniform distribution in the raw material. In the process of storing control and experimental samples in a cooled state, the intensity of accumulation of oxidation products in them was studied weekly according to acid and peroxide numbers by common methods. The results are processed by methods of mathematical statistics.

The second series of studies was to confirm the effectiveness of the antioxidant action of dihydroquercetin during the long-term storage of the mechanically deboned meat in a frozen state. Mechanical boned broiler meat contains skin that, due to its fat content, is significantly susceptible to oxidative spoilage; therefore, it is important to reduce the degree of its influence on the formation of oxidation products in raw materials [2].

The preparation was added to the experimental samples of mechanically deboned meat at a dose of 0.50; 0.75; and 1.00 kg per 100 kg of raw material, in hydrated form. During 6 months of storage at a temperature of -18 ± 1 °C, a monthly determination of the content of oxidation products — acid and peroxide numbers was also carried out in 3-fold repetition in the control and experimental samples.

In accordance with the aim and objectives of the study, the objects of the research were:

- as antioxidants - “Dihydroquercetin” (DHQ), “Ascorbic acid”, “Tocopherol” and “Rutin” [7,8,9,10];
- fillet of chilled carcasses of broiler chickens with pH24 6.2 – 6.5, according to GOST R 52702-2006 [13];
- mechanically deboned meat of broiler chickens according to GOST 31490-2012 [14];
- skin from carcasses of broiler chickens.

Research results. The issue of storage meat products is especially relevant in poultry farming [11] and in beef breeding [12] in view of the wide range of products for the mass consumer.

The comparative analysis and comprehensive assessment of the content of oxidation products in the samples fairly testify to the effect of antioxidants on the reduction of oxidative spoilage of raw materials, but with different efficiencies (table 2).

Table 2 – Change in the acid number of samples, mg KOH/g

Raw material	Control	Vitamin E	Vitamin C	Rutin	DHQ
<i>At the beginning of the experiment:</i>					
Skin	0.196±0.020				
MDM	0.178±0.030				
Fillet	0.112±0.040				
<i>After 7 days of storage:</i>					
Skin	0.936±0.03	0.248±0.01	0.233±0.01	0.218±0.01	0.210±0.01
MDM	0.798±0.24	0.247±0.02	0.223±0.02	0.195±0.01	0.180±0.01
Fillet	0.668±0.57	0.134±0.04	0.122±0.03	0.116±0.04	0.112±0.04
<i>After 14 days of storage:</i>					
Skin	1.028±0.03	0.820±0.01	0.743±0.03	0.603±0.01	0.530±0.01
MDM	0.839±0.10	0.572±0.44	0.443±0.02	0.395±0.01	0.348±0.07
Fillet	0.719±0.01	0.554±0.01	0.422±0.03	0.321±0.03	0.262±0.03
<i>After 21 days of storage:</i>					
Skin	2.153±0.04	1.895±0.02	1.782±0.01	1.691±0.06	1.619±0.08
MDM	1.985±0.10	1.836±0.15	1.652±0.03	1.616±0.03	1.524±0.03
Fillet	1.974±0.09	1.793±0.04	1.593±0.07	1.543±0.07	1.438±0.02
<i>After 28 days of storage:</i>					
Skin	2.542±0.02	2.389±0.01	2.125±0.07	1.925±0.04	1.705±0.04
MDM	2.306±0.06	2.194±0.05	1.897±0.03	1.803±0.04	1.694±0.03
Fillet	2.200±0.08	2.122±0.02	1.822±0.04	1.794±0.06	1.525±0.03

When studying the antioxidant activity of preparations, in parallel with the acid number, the peroxide number was determined - an indicator that characterizes the amount of primary lipid oxidation products (hydroperoxides and peroxides) in experimental samples of raw materials (table 3).

Table 3 – Change in the peroxide number of samples, mmol (SO₂)/kg

Raw material	Control	Vitamin E	Vitamin C	Rutin	DHQ
<i>At the beginning of the experiment:</i>					
Skin	0.008±0.0008				
MDM	0.007±0.0012				
Fillet	0.004±0.0004				
<i>After 7 days of storage:</i>					
Skin	2.346±0.20	0.131±0.02	0.098±0.01	0.063±0.01	0.033±0.01
MDM	2.138±0.22	0.118±0.02	0.086±0.01	0.043±0.01	0.025±0.01
Fillet	1.154±0.09	0.078±0.01	0.033±0.01	0.025±0.01	0.013±0.01
<i>After 14 days of storage:</i>					
Skin	3.459±0.09	2.911±0.12	1.443±0.06	0.903±0.08	0.473±0.08
MDM	3.038±0.72	2.708±0.59	1.141±0.34	0.808±0.08	0.360±0.19
Fillet	2.765±0.06	2.078±0.01	1.033±0.01	0.715±0.01	0.313±0.14
<i>After 21 days of storage:</i>					
Skin	4.422±0.08	4.192±0.10	3.948±0.07	3.822±0.05	3.719±0.04
MDM	4.387±0.09	4.106±0.52	3.903±0.03	3.721±0.06	3.659±0.15
Fillet	4.072±0.09	3.922±0.01	3.872±0.08	3.715±0.02	3.595±0.09
<i>After 28 days of storage:</i>					
Skin	5.620±0.03	4.368±0.09	4.232±0.03	4.152±0.09	3.870±0.02
MDM	5.494±0.09	4.284±0.06	4.162±0.03	4.091±0.03	3.829±0.05
Fillet	5.452±0.04	4.219±0.05	4.105±0.08	3.986±0.06	3.729±0.07

In the control skin sample from broiler carcasses, after 7 days of storage of raw materials, the peroxide value reached 2.3459 mmol (SO₂)/kg, which characterizes the sample by this indicator as fresh, but not subject to storage. The introduction of antioxidants into experimental samples of raw materials inhibits the formation of lipid oxidation products, therefore, the peroxide number in them was significantly less.

So, with the addition of vitamin E, the peroxide value in the experimental samples was 1.3 times lower than in the control ones. With the introduction of vitamin C, rutin and dihydroquercetin as antioxidants, this indicator in the samples of raw materials was even lower - respectively 1.6; 1.7 and 1.9 times.

The introduction of antioxidants into experimental samples of raw materials in the studied concentrations ensured within 28 days of storage, a lower value of the indicators of oxidative spoilage: the acid number by an average of 74%, and the peroxide number by 1.5-1.7 times, relative to these indicators in the control sample.

To confirm the effectiveness of the antioxidant action of dihydroquercetin during long-term storage of mechanically deboned meat, the second series of studies was held.

The comparative analysis and comprehensive assessment of the accumulation of oxidation products in the control and experimental samples during long-term storage indicates a significant inhibitory effect of dihydroquercetin on the rate of formation of substances that cause oxidative spoilage of mechanically deboned meat of broiler chickens.

The acid number indicates the formation of free fatty acids in raw materials resulting from hydrolytic spoilage of fat. In the studied meat samples, after 1 month of storage, this indicator had the highest value in the control sample that did not contain an antioxidant preparation (0.3873 mg KOH/g). After 6 months of storage, the acid number increased and reached 0.3896 mg KOH/g, which exceeded this indicator in

experimental samples of raw materials at dihydroquercetin levels, respectively: 0.50 kg/100 kg - by 0.0071 mg KOH/g; 0.75 kg/100 kg - per 0.0253 mg KOH/g and at preparation concentration of 1.0 kg/100 kg - 0.0368 mg KOH/g (table 4).

Table 4 – Change in the acid number of mechanically deboned meat during storage (mg KOH/g)

Shelf life	Control	Dihydroquercetin concentration, kg/100 kg		
		0.50	0.75	1.00
1 month	0.3873±0.0199	0.3799±0.0122	0.3624±0.0129	0.3470±0.0093
2 month	0.3876±0.0201	0.3800±0.0120	0.3626±0.0193	0.3473±0.0094
3 month	0.3878±0.0157	0.3803±0.0207	0.3629±0.0162	0.3476±0.0017
4 month	0.3881±0.0230	0.3807±0.0143	0.3633±0.0134	0.3482±0.0192
5 month	0.3889±0.0210	0.3812±0.0621	0.3638±0.0267	0.3495±0.0815
6 month	0.3896±0.0308	0.3825±0.0932	0.3643±0.0384	0.3528±0.0631

While testing, it was established that the antioxidant activity of dihydroquercetin is also effective in the formation of primary lipid oxidation products characterized by the peroxide number (table 5).

Table 5 – Change in the peroxide number of mechanically deboned meat during storage (ммоль(1/2O₂)/кг)

Shelf life	Control	Dihydroquercetin concentration, kg/100 kg		
		0.50	0.75	1.00
1 month	0.0083±0.0008	0.0066±0.0018	0.0058±0.0092	0.0049±0.0012
2 month	0.0084±0.0027	0.0068±0.0046	0.0060±0.0005	0.0052±0.0032
3 month	0.0086±0.0078	0.0069±0.0026	0.0061±0.0015	0.0055±0.0063
4 month	0.0089±0.0088	0.0071±0.0226	0.0063±0.0505	0.0057±0.0612
5 month	0.0092±0.0042	0.0074±0.0066	0.0065±0.0071	0.0060±0.0045
6 month	0.0095±0.0065	0.0077±0.0417	0.0068±0.0731	0.0063±0.0921

The introduction of the dihydroquercetin antioxidant in the composition of the experimental samples in the studied concentrations ensured, after 1 and 6 months of storage, a lower value of the indicators of oxidative spoilage: the acid number by 10%, and the peroxide number by 1.5-1.7 times, relative to these indicators in the control sample.

**Н. Н. Кузьмина¹, О. Ю. Петров¹, В. Г. Семенов²,
Д. А. Баймұқанов³, А. И. Любимов⁴, Х. А. Аубакиров⁵**

¹Марийск мемлекеттік университеті, Йошкар-Ола, Марий Эл Республикасы, Ресей;

²Чуваш мемлекеттік ауылшаруашылық академиясы, Чебоксары, Чуваш Республикасы, Ресей;

³«Қазақ ұлттық аграрлық университеті» коммерциялық емес акционерлік қоғамы, Алматы, Қазақстан;

⁴Ижевск мемлекеттік ауылшаруашылық академиясы, Ижевск, Удмурт Республикасы, Ресей;

⁵М. Х. Дулати атындағы Тараз мемлекеттік университеті, Тараз, Қазақстан

ҚҰС ӨНДЕУ ШИКІЗАТЫН САҚТАУДА ТАБИҒИ АНТИОКСИДАНТТАРДЫҢ ТИІМДІЛІГІН ЗЕРТТЕУ

Аннотация. Өңделген құс шикізатының сақтау мерзімін ұлғайту үшін антиоксидант ретінде дигидрокверцетинді қолданудың тиімділігі зерттелген. Табиғи антиоксидантты дәрумендер Е, С, рутин және дигидрокверцетиннің шайналған ет, механикалық өңделген ет және терісімен бірге балапан – бройлер ұшасының сақталуына ұтымды әсер етудің салыстырмалы бағалауы жасалды. Тәжірибиелі үлгілерге антиоксидантарды енгізу шикізатқа 28 тәулік бойы тотығу көрсеткішінің аз мәнін, қышқылдану 74%, ал ашу санын 1,5-1,7 есе төмендетті.

Алынған нәтижелер дигидрохверцетинді аналогтарымен қатар, қолданудың жоғары тиімді екендігіне көз жеткізді.

Үлгілердегі тотығу өнімдері мөлшерінің салыстырмалы талдауы және комплексті бағалауы үлгілерде антиоксидат әсерінен шикізаттың тотыға бұзылуына әсер ететіндігін көрсетеді, алайда түрлі тиімділікпен жүзеге асырылады.

Антиоксидантарды бақылау үлгідегі шикізатқа қосу оның айтарлықтай тотыға бүліну үдерісін тоқтатады. Е дәруменін ендіргенде 28 тәулік сақтау барысында қышқылдану саны 0,84 мг КОН/г орташа төмен болады, С дәруменін, рутин және дигидрохверцетинді сәйкесінше бақылау үлгімен салыстырғанда 1,27; 3,52 және 3,61 мг КОН/г көрсетеді.

Тәжірибиелі үлгілердің құрамына дигидрохверцетин қосқанда оның тотыға бүліну үдерісін тоқтатады. Антиоксидантты негізгі шикізатқа 0,5 кг/100 кг деңгейінде қосу, бір ай сақтаудан кейін бақылаумен салыстырғанда, қышқылдану 2%, сәйкесінше 0,75 және 1,0 кг – 6 және 10 %, төмендейді. Тәжірибе барысында және 6 ай сақтаудан кейін үлгілерде бақылаумен салыстырғанда сәйкесінше 2, 6 и 9 % төмен көрсетті.

Бақылау және тәжірибе үлгілеріндегі тотығу өнімдерінің жинақталуын салыстырмалы талдау және комплексті бағалау дигидрохверцетиннің механикалық өңделген балапан – бройлерлерді ұзақ сақтау барысында тотыға бүлінуді түзетін заттардың тежелуіне айтарлықтай әсер етеді.

Алынған мәліметтер дигидрохверцетин, басқа табиғи антиоксиданттармен салыстырғанда, өңделген құс шикізатында тотығу өнімдерінің түзілуін тиімді тежейді. Атап айтқанда, ірі қара ет өндірісінде келесідей жұмыстарды жүзеге асыруға мүмкіндік береді: сақтау мерзімі ұзақ өнімдерді өндіру; ағзадағы бос – радикалды үдерістерді тежейтін антиоксиданттармен қамтамасыз ету; функционалды бағыттағы азық өнімдерінің ассортиментін кеңейту.

Өңделген құс өнімдерінің сақтау мерзімін ұзартуда дигидрохверцетиннің антиоксиданттық белсенділігі жоғары екендігі анықталды, тотығу өнімдерінің жинақталуын тежейді.

Түйін сөздер: құс өңдеу, балапан-бройлер ұшаларының терісі, механикалық етті сүйектен айыру, балапан-бройлердің салқындалатын сүбесі, табиғи антиоксидант, Е, С дәрумені, рутин, дигидрохверцетин, тотығу өнімдері, сақтау ұзақтығы.

**Н. Н. Кузьмина¹, О. Ю. Петров¹, В. Г. Семенов²,
Д. А. Баймуканов³, А. И. Любимов⁴, Х. А. Аубакиров⁵**

¹Марийский государственный университет, Йошкар-Ола, Республика Марий Эл, Россия;

²Чувашская государственная сельскохозяйственная академия,

Чебоксары, Чувашская Республика, Россия;

³Некоммерческое акционерное общество

«Казахский национальный аграрный университет», Алматы, Казахстан;

⁴Ижевская государственная сельскохозяйственная академия,

Ижевск, Удмуртская республика, Россия;

⁵Таразский Государственный университет им. М.Х. Дулати, Тараз, Казахстан

ИССЛЕДОВАНИЕ ЭФФЕКТИВНОСТИ ПРИРОДНЫХ АНТИОКСИДАНТОВ ПРИ ХРАНЕНИИ СЫРЬЯ ПТИЦЕПЕРЕРАБОТКИ

Аннотация. Исследованы эффективность использования дигидрохверцетина в качестве антиоксиданта, способствующего увеличению продолжительности хранения сырья птицепереработки.

Проведена сравнительная оценка эффективности действия природных антиоксидантов витаминов Е, С, рутина и дигидрохверцетина на продолжительность хранения образцов фаршей из филе, мяса механической обвалки и кожи с тушек цыплят-бройлеров. Введение антиоксидантов в опытные образцы сырья обеспечило в течение 28 дней хранения меньшее значение показателей окислительной порчи: кислотного числа в среднем на 74 %, а перекисного числа – в 1,5–1,7 раза, относительно этих показателей в контрольном образце без добавления антиоксидантов.

Применение дигидрохверцетина в опытных образцах замороженного мяса механической обвалки для длительного хранения в исследуемых концентрациях обеспечило уже через 1 месяц хранения меньшее значение показателей окислительной порчи: кислотного числа на 2–10 %, а перекисного числа – в 1,5 и 1,7 раза, относительно этих показателей в контрольном образце.

Полученные результаты убеждают в высокой эффективности применения дигидрохверцетина в качестве активного антиоксиданта, что обеспечивает возможность его использования наряду с имеющимися аналогами.

Сравнительный анализ и комплексная оценка содержания продуктов окисления в образцах объективно свидетельствуют о влиянии антиоксидантов на снижение окислительной порчи сырья, но с разной эффективностью.

Добавление антиоксидантов в контрольные образцы сырья способствовало существенному торможению его окислительной порчи. При введении витамина Е, кислотное число за 28 дней хранения оказалось ниже в среднем на 0,84 мг КОН/г, при добавлении витамина С, рутина и дигидрокверцетина, соответственно, на 1,27; 3,52 и 3,61 мг КОН/г - по сравнению с контрольным образцом.

Добавление дигидрокверцетина в состав опытных образцов способствовало торможению его окислительной порчи. При введении антиоксиданта на уровне 0,5 кг/100 кг основного сырья после 1 месяца хранения кислотное число оказалось меньше, чем в контрольном образце на 2 %, при добавлении 0,75 и 1,0 кг – на 6 и 10 %, соответственно. Подобная закономерность прослеживалась в течение всего периода эксперимента и после 6 месяцев хранения, значения этого показателя в опытных образцах оказались меньше контроля на 2, 6 и 9 %, соответственно.

Полученные результаты свидетельствуют, что дигидрокверцетин, по сравнению с другими природными антиоксидантами, наиболее эффективно ингибирует образование продуктов окисления в сырье птицепереработки, что обуславливает возможность его применения при производстве мясных продуктов, в частности в мясном скотоводстве, и создает условия для: производства продуктов с пролонгированным сроком годности; обеспечения организма антиоксидантами, предотвращая свободно-радикальные процессы в нем; расширения ассортимента продуктов питания функциональной направленности.

Объективно установлено, что дигидрокверцетин проявляет высокую антиоксидантную активность, препятствуя накоплению продуктов окисления, и способствует значительному увеличению сроков хранения сырья птицепереработки.

Ключевые слова: птицепереработка, кожа с тушек цыплят-бройлеров, мясо механической обвалки, филе охлажденных тушек цыплят-бройлеров, природные антиоксиданты, витамины Е, С, рутин, дигидрокверцетин, продукты окисления, продолжительность хранения.

Information about the authors:

Kuzmina Nadezhda Nikolayevna, Assistant Faculty, Department of Technology of Meat and Dairy Products, Mari State University, Yoshkar-Ola, Mari El Republic, Russia; kuzmina221995@mail.ru; <https://orcid.org/0000-0002-8552-6364>

Petrov Oleg Yuryevich, Doctor of Agricultural Sciences, Associate Professor of the Department of Technology of Meat and Dairy Products, Mari State University, Yoshkar-Ola, Mari El Republic, Russia; tmspetrov@yandex.ru; <https://orcid.org/0000-0002-3647-5345>

Semenov Vladimir Grigorievich, Doctor of Biological Sciences, Professor, Honored Scientist of the Chuvash Republic, Professor of the Department of Morphology, Obstetrics and Therapy of the Chuvash State Agricultural Academy, Cheboksary, Chuvash Republic, Russia; semenov_v.g@list.ru; <https://orcid.org/0000-0002-0349-5825>

Baimukanov Dastanbek Asylbekovich, Corresponding Member of the National Academy of Sciences of the Republic of Kazakhstan, Doctor of Agricultural Sciences, Professor of the Department of Physiology, Morphology, and Biochemistry named after academician N.U. Bazanova, NJSC “Kazakh National Agrarian University”, Almaty, Kazakhstan; dbaimukanov@mail.ru; <https://orcid.org/0000-0002-4684-7114>

Lyubimov Alexander Ivanovich, Doctor of Agricultural Sciences, Professor, Honorary Worker of the AIC of Russia, Honored Worker of Agriculture in the Russian Federation, Head of the Scientific School “Development of Breeding and Technological Methods of Intensification of Livestock Production”, Head of the Laboratory “Breeding, Reproduction and Prevention of Diseases of Farm Animals”, Rector of the Izhevsk State Agricultural Academy, Izhevsk, Udmurt Republic, Russia; rector@izhgsha.ru; <https://orcid.org/0000-0001-5369-6249>

Aubakirov Khamit Abilgazievich, Candidate of Agricultural Sciences, Associate Professor of the Department of Biotechnology, M.H. Dulati Taraz State University, Taraz, Kazakhstan; hamit_a57@mail.ru; <https://orcid.org/0000-0003-2670-4834>

REFERENCES

- [1] Denisovich Yu.Yu., Borozda A.V., Mandro N.M. (2012) Development of technology of enriched meat products of a functional orientation // Bulletin of the Altai State Agrarian University. Barnaul. N 6 (92). P. 83-87 (in Russ.).
- [2] Antipova L.V., Glotova I.A. (2006) The use of secondary collagen-containing raw materials of the meat industry. SPb. GIORD. 384 p. (in Russ.).
- [3] Mandro N.M., Borozda A.V., Denisovich Yu.Yu. (2009) Development of technology for minced meat using a natural antioxidant // Bulletin of the Altai State Agrarian University. Barnaul. N 5 (55). P. 72-75 (in Russ.).
- [4] Aniksevich O.N. (2011) Dihydroquercetin in the meat processing industry. Food industry: science and technology. N 3 (13). P. 38-42 (in Russ.).
- [5] Borozda A.V., Denisovich Yu.Yu. (2009) New aspects in the use of dihydroquercetin in the production of semi-finished meat products. Agrarian science - to agriculture. Proceedings of IV International scientific - practice. conf. Barnaul. P. 25-27 (in Russ.).
- [6] Recommended levels of consumption of food and biologically active substances (2004) Methodological recommendations of the State Sanitary and Epidemiological Regulation of the Russian Federation N 2.3.1.1915-04 of 2004 (in Russ.).
- [7] SanPiN 2.3.2.1078-2011 "Hygienic requirements for safety and nutritional value of food products (as amended on July 6, 2011)" (in Russ.).
- [8] TR 9100-241-21428156-2011 "Dihydroquercetin biologically active food supplement".
- [9] MP-003326 dated 23.11.2015 Alpha-Tocopherol Acetate (Vitamin E).
- [10] MP-000043 dated 13.04.07 Ascorutin.
- [11] Alekseev I.A., Semenov V.G., Baimukanov D.A., Alekseev V.A., Evdokimov N.V., Yakimov A.V., Obukhova A.V. (2019). Basulifor probiotic supplement, its impact on body and productivity of young quails // Bulletin of the National Academy of Sciences of the Republic of Kazakhstan. Vol. 4, N 380 (2019), 27-37. ISSN 2518-1467 (Online), ISSN 1991-3494 (Print). <https://doi.org/10.32014/2019.2518-1467.89>
- [12] Baimukanov D.A., Pristupa V.N., Kolosov Yu.A., Donnik I.M., Torosyan D.S., Kolosov A.Yu., Orlova O.N., Yuldashbayev Yu.A., Chylbak-ool S.O. (2019) Improvement of breeding and productive traits of Kalmyk cattle breed // Bulletin of the National Academy of Sciences of the Republic of Kazakhstan. Vol. 2, N 378 (2019), 128-145. ISSN 2518-1467 (Online), ISSN 1991-3494 (Print). <https://doi.org/10.32014/2019.2518-1467.51>
- [13] GOST R 52702-2006 "Chicken meat (carcasses of chickens, broiler chickens, and parts). Technical conditions (in Russ.).
- [14] GOST 31490-2012 "Mechanically deboned poultry meat. Technical conditions.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1467 (Online), ISSN 1991-3494 (Print)

<http://www.bulletin-science.kz/index.php/en/>

Редакторы *М. С. Ахметова, Д. С. Аленов, А. Ахметова*
Верстка на компьютере *Д. А. Абдрахимовой*

Подписано в печать 14.08.2020.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
21,6 п.л. Тираж 500. Заказ 4.